## ADVANCED CYSYLLTIADAU CONNECTIONS PELLACH SESSION PLAN

#### **Physics Tutorial 2: Performing Like a Physicist**

#### <u>Aim</u>

To understand what standard form is, and how it is used to express very large or small values in physics

#### <u>Learning Objectives</u>



By the end of this session, the mentees will be able to...

- Explain when and why physicists use standard form
- Perform basic calculations using standard form
- Recognise the difference between accuracy and precision, and how they are used in physics

#### **Prepare in Advance**

• There shouldn't be anything to prepare in advance, just have a read through the answers and make sure you understand and can explain them!

#### <u>Scaffolding</u>

If a mentee finishes early:

- Get them to try the optional extras (if they haven't already)
- Get them to think of another "million pound question" about a physics topic of their choice

If a mentee is struggling to calculate using standard form:

 Talk them through it stepby-step (e.g. "are you multiplying or dividing?" "what is the rule you need to use?"

If a mentee is struggling to think of a "million pound question":

- Allow them to choose any physics topic instead of just from this session
- Show them some example questions





## MENTOR GUIDANCE SESSION 2: PERFORMING LIKE A PHYSICIST PART 1: SELF-STUDY RECAP

This first half of this tutorial is going through the self-study materials to ensure that mentees understand what standard form is, why we use it, and how to calculate with it.

# **TASK 1: CALCULATION PRACTICE**

1. $(3 \times 10^4) \times (8 \times 10^2)$ $= (3 \times 8) \times (10^4 \times 10^2)$ $= 24 \times 10^{4+2}$ Mentees may forget to $= 24 \times 10^6$ between 0 and 10 $= 2.4 \times 10^7$ (8.4 × 10 <sup>6</sup> )	$\begin{array}{l} \textbf{4.} & \left(5 \times 10^{6}\right) \times \frac{2.4 \times 10^{2}}{1.2 \times 10^{8}} = 10 \\ & = \left(5 \times 10^{6}\right) \times \left(\left(2.4 \div 1.2\right) \times \left(10^{2} \div 10^{8}\right)\right) \\ & = \left(5 \times 10^{6}\right) \times \left(2 \times 10^{2-8}\right) \\ & = \left(5 \times 10^{6}\right) \times \left(2 \times 10^{-6}\right) \\ & = \left(5 \times 2\right) \times \left(10^{6} \times 10^{-6}\right) \end{array}$
$ \begin{array}{l} \hline \hline (2.1 \times 10^2) \\ = (8.4 \div 2.1) \times (10^6 \div 10^2) \\ = 4 \times 10^{6-2} \\ = 4 \times 10^4 \end{array} $	$=10 imes 10^{6-6} \ =10 imes 10^{0} \ =10$
$\begin{array}{l} \textbf{3.} \left(9 \times 10^2\right) \times \frac{\left(5 \times 10^9\right)}{\left(2.5 \times 10^3\right)} \\ &= \left(9 \times 10^2\right) \times \left((5 \div 2.5) \times \left(10^9 \div 10^3\right)\right) \\ &= \left(9 \times 10^2\right) \times \left(2 \times 10^{9-3}\right) \\ &= \left(9 \times 10^2\right) \times \left(2 \times 10^6\right) \\ &= \left(9 \times 2\right) \times \left(10^2 \times 10^6\right) \\ &= 18 \times 10^{2+6} \\ &= 18 \times 10^8 \\ &= 1.8 \times 10^9 \end{array}$	5. $\frac{10^{3} \times 10^{-2} \times 10^{7}}{10^{-5} \times 10^{4}} = 10^{9}$ $= (10^{3-2+7}) \div (10^{-5+4})$ $= (10^{8}) \div (10^{-1})$ $= 10^{81}$ Mentees may not have tried the harder ones (4 and 5)

### **COMMON PROBLEMS**

Mentees might:

- Forget that the value has to be between 0 and 10, so may not convert (e.g. from 24 to 2.4) they might also forget to change the power when they do
- Be confused by the different format in Q4-5 ( $10^{B}$  instead of A x  $10^{B}$ ). You can explain that this is the same as writing 1 x  $10^{B}$  in a shorter form



## MENTOR GUIDANCE **SESSION 2: PERFORMING LIKE A PHYSICIST** PART 1: SELF-STUDY RECAP

## **TASK 2: GRAVITY**

There are **two correct methods** to solve this. We've included both here:

$$egin{aligned} F &= rac{Gm_1m_2}{r^2} &= ((6.67 imes 10^{-11}) imes (5.97 imes 10^{24}) imes (7.34 imes 10^{22})) \div (3.85 imes 10^8) \ &= ((6.67 imes 5.97 imes 7.34) imes (10^{-11} imes 10^{24} imes 10^{22}) \div (3.85 imes 10^8) \ &= (292.28 imes 10^{35}) \div (3.85 imes 10^8) \ &= (2.92 imes 10^{37}) \div (3.85 imes 10^8) \ &= (292.28 \div 3.85) imes (10^{35} \div 10^8) \ &= (2.92 \div 3.85) imes (10^{37} \div 10^8) \ &= 75.9 imes 10^{27} \ &= 7.59 imes 10^{28} \ &= 7.59 imes 10^{28} \end{aligned}$$

# **TASK 3: ACCURACY VS PRECISION**

The idea of this exercise is to get mentees to consider what accuracy and precision are in the context of a real experiment. Mentees might suggest the following things:



• A radio-controlled clock is accurate to the actual time



- Measuring to the nearest 100 ms makes the experiment more precise
- The physicist is performing the experiment every day, so if they do it lots of times their measurements will be precise



• Human error of the person taking the time is not very accurate

 $G = 6.67 \times 10^{-11} \,\mathrm{Nm^2/kg^2}$ 

 $m_1 = 5.97 \times 10^{24} \, {
m kg}$  $m_2 = 7.34 \times 10^{22} \, \mathrm{kg}$ 

 $r = 3.85 \times 10^8 \text{m}$ 

- The horizon being obscured makes the dawn time measured less accurate
- What way is their window facing? This will make the measurement less accurate to actual dawn time



## LOW PRECISION

• Human error also makes the time less precise - they are measuring when the light reaches their room and taking the time by eye



## MENTOR GUIDANCE SESSION 2: PERFORMING LIKE A PHYSICIST PART 1: SELF-STUDY RECAP

## **TASK 4: RECORDING DATA**

Mentees are asked to find **at least 3**, but there are quite a few errors in the table. They're all pointed out here:



# OPTIONAL EXTRAS 2. $E = \frac{Q}{4\pi\varepsilon_0 r^2}$

**1.**  $E = mc^2$  $= \left(3.20 imes 10^{-16}
ight) \div \left(4 \pi imes \left(8.85 imes 10^{-12}
ight) imes \left(2.4 imes 10^{-9}
ight)^2
ight)$  $= ig( 9.11 imes 10^{-31} ig) imes ig( 2.99 imes 10^8 ig)^2$  $= (3.20 imes 10^{-16}) \div (4\pi imes (8.85 imes 10^{-12}) imes (2.4 imes 10^{-18}))$  $=(9.11 imes 10^{-31}) imes (2.99 imes 10^{16})$  $= (3.20 imes 10^{-16}) \div (4\pi imes (8.85 imes 2.4) imes (10^{-12-18}))$  $= (9.11 imes 2.99) imes (10^{-31} imes 10^{16})$  $= (3.20 imes 10^{-16}) \div (4\pi imes (21.24 imes 10^{-30}))$  $=(3.20 imes 10^{-16}) \div (266.91 imes 10^{-30})$  $= 27.24 imes 10^{-15}$  $= (3.20 \div 266.91) imes \left( 10^{-16} \div 10^{-30} 
ight)$  $=2.72 imes10^{-14}$  $m_e = 9.11 \times 10^{-31}$ kg  $= 0.0120 imes 10^{14}$  $c = 2.99 \times 10^8$  m/s  $=1.20 imes10^{12}$ 

 $Q = 3.20 \times 10^{-16} C$ 

 $arepsilon_0 =$  8.85 × 10<sup>-12</sup> F/m r = 2.4 nm



## MENTOR GUIDANCE SESSION 2: PERFORMING LIKE A PHYSICIST PART 2: SKILLS PRACTICE

The second half of this tutorial gives students an opportunity to practice their skills and take part in a quiz.

# WHO WANTS TO BE A MILLIONAIRE?

In 2001, a man cheated his way to winning Who Wants to be a Millionaire: <a href="https://www.youtube.com/watch?v=HIGtLRnGCD4">https://www.youtube.com/watch?v=HIGtLRnGCD4</a> (5 mins)

The million pound question was...

A number one followed by one hundred zeros is known by what name?



**Your mentees would have got the million pound question correct!** They know that mega has 6 zeroes, giga has 9 zeroes, and nano has 9 zeroes after the decimal place. So it's got to be a Googol.

Give the mentees 5-10 minutes to **come up with their own "million pound question"** about this session's topics. Mentees can work in pairs/threes in the mini breakout rooms, or on their own. **Make sure they create 1 question with 4 multiple choice answers.** 

After the time is up, bring mentees back to the main room and ask them to **privately** message you their question and choices in the Zoom chat (don't forget to get the correct answer from them too!).

Put the questions into the template on the PPT, then quiz the mentees and see how many they can get correct.

## TOP TIPS

- If mentees complete their question early, you can add their questions as you go to save time.
- Mentees might get stuck. Encourage them to work together, or use some of the examples on the PPT to give them inspiration.
  - If they're really stuck, you could get them to make a question about any physics topic of their choice.

You could add a competitive element with 1st, 2nd and 3rd places!



#### **PHYSICS TUTORIAL 2: PERFORMING LIKE A PHYSICIST**

# **BRIGHT IDEAS!**

This page contains ideas for alternative sessions, changes/additions, extra activities, etc. Feel free to use as you wish!

## Standard Form Everywhere

You could include examples of where standard form is useful in various careers and situations. Some examples:

- Engineering and construction: large structures, bridges, roads
- Architecture: maths is used throughout architecture
- Biology and chemistry: microscopic level, molecules
- Finances: large companies, revenue
- Physics (of course!): astronomy, semiconductors

<sup>5</sup> Did you know that engineering has it's own standard form notation called "engineering form"?



# Physics Skills

If you think the mentees are struggling to link the topics to wider skill applications, you can run an activity where they list the skills they use to solve the problems, and then discuss where they use those skills in physics.

## Star Wars Standard Form

If mentees are struggling with standard form calculations, try this activity.

The Millenium Falcon flies at the speed of sound normally, and flies at the speed of light when it goes into hyperdrive.

Calculate how long these journeys would take the Millenium Falcon (answers in standard form!)

speed of sound =  $3.40 \times 10^2$  m/s speed of light =  $2.99 \times 10^8$  m/s

From	То	Distance (km)	Hyper drive?	Time Taken (s)
Hoth	Naboo	$1.224 \times 10^{7}$	NO	??
Tatooine	Dagobath	3.06 × 10 <sup>6</sup>	NO	??
Endor	Ord Mantell	1.35 × 10 <sup>13</sup>	YES	??
Dantooine	Kashyyk	??	NO	$1.2 \times 10^{4}$
Bespin	Kessei	??	YES	$7.5 \times 10^{3}$

MENTORING PROJECT